

Conifer Encroachment

Sage grouse breeding habitat is lost to invading junipers when trees exceed 4% canopy cover. Focused tree removal prevents conversion of sagebrush steppe to conifer woodlands and benefits sage grouse populations.

Benefits of conifer control

- Maintaining native understory plants.
- 2. Reducing risk of large and severe wildfires.
- 3. Improving habitat for declining species.
- 4 Reducing soil erosion and conserving soil water.
 - 5 Increasing ecosystem resilience to fire and resistance to cheatgrass.

know that trees catch the snow? Removing encroaching conifer stands from sagebrush ecosystems helps keep snow on the ground longer during the spring. This allows water to seep slowly into the ground to better sustain plants, streams, and wildlife during the West's hot, dry summer.



nrcs.usda.gov/wildlife sagegrouseinitiative.com

MIDIFF

Conservation measures set in place for sage grouse significantly increase the protection for mule deer migration and winter habitat.

Easements protect large working lands from subdivision development in sage

agreements that provide a

and preserve habitat.

grouse core areas by using voluntary

financial boost for ranchers

The Sage Grouse Initiative is a partnership-based, science-driven effort that uses voluntary incentives to proactively conserve America's western rangelands, wildlife, and rural way of life. This initiative is part of Working Lands For Wildlife, which is led by USDA's Natural Resources Conservation Service.

SHE GROSE IS AN UMBLE Studies have shown that conservation efforts for sage grouse also help 350 other species of plants, invertebrates, amphibians, reptiles, birds and

conservation efforts for sage grouse also help **350** other species of plants, invertebrates, amphibians, reptiles, birds and mammals inhabiting the sagebrush sea, like mule deer and songbirds.

Wet "mesic"

habitat, or places

where water meets

land comprise less than 2% of the

entire landscape in the

West. Protecting and

restoring these "emerald

islands" in the desert

benefits livestock ranching and

wildlife, including sage grouse.

Songbirds like the Brewer's sparrow, sagebrush sparrow and sage thrasher are 13-19% more abundant near large leks,

Sage grouse can collide with livestock fences when flying low over sagebrush. When grouse fly into their breeding grounds, or leks," the dim predam light makes it dissignit to avoid foing some seasons.

"leks," the dim predawn light makes it difficult to avoid foing vires. By placing made

wires. By placing markers on fence wires, ranchers as \$3%.

their part to reduce the potential for fence collision by

which support half of all sage grouse populations.

Raptors

Conifers provide perches for predators.

Invasive Grasses

The spread of invasive annual grasses are highly flammable and dry out earlier than native plants, leading to more frequent, hotter fires. Once sagebrush habitat burns in a megafire, it's hard to restore, leaving noxious weeds that degrade rangelands and wildlife habitat.

the sagebrush sea's diverse, wildflowers, and bunchgrasses That's bad for birds and herds, which rely on nutritious, native perennial plants.

grasses, like cheatgrass and medusahead, are linked to unwanted wildfires. Invasive

> These invasive grasses replace native plants - like sagebrush, - with a monoculture of weeds.

Pest rotation grazing boosts the abundance of arthropods, like spiders and for nutrition pastures from grazing host the most anthropods important for nutrition.

Conservation on working western landscapes restores sagebrush communities where sage grouse and other wildlife share the same extensive home range, and helps ranchers pass on their legacy and rural way of life.

